

The Long QT Syndrome

A Guide for Patients and Health Care Providers



Why do I need to know about the Inherited Long QT syndrome?

We estimate that 1 in 2,000 people have LQTS. LQTS-precipitated sudden deaths continue to claim otherwise healthy infants, children, adolescents, and adults at an unacceptably high rate. However, with increased awareness, genetic testing, and effective treatment options, LQTS can be diagnosed early and sudden death prevented. Still, this condition is often undetected prior to death and not recognized as the cause of death. Family members of individuals with unexplained death should be tested for LQTS and other genetic arrhythmias. If an inherited condition like LQTS was responsible for the unexplained death, then 50% of family members may also be affected. LQTS is a treatable disorder and, with correct diagnosis and treatment, most deaths are preventable. Currently mortality for properly treated patients is close to 1%.

Physicians need to know:

- When to consider LQTS as a possible diagnosis.
- When to refer patients for diagnosis & treatment.
- About genetic testing for LQTS and other SADS conditions.
- How to develop a family pedigree and screen family members for LQTS.

What is LQTS?

LQTS is a disturbance of the heart's electrical system. It is caused by abnormalities of microscopic pores (proteins) in the heart cells called ion channels. Ions such as potassium, sodium, calcium and chloride pass back and forth across the cell membrane through ion channels. As they do, these ion channels generate the electrical activity (depolarization and repolarization) that controls the heart's beating. Our window to this electrical activity of the heart is through an electrocardiogram (EKG or ECG). Potassium and sodium ion channels are two classes of ion channels affected in LQTS. The abnormal channels prolong the repolarization ("recharging") process and the QT interval, thus predisposing patients to certain cardiac arrhythmias. Thus, LQTS is a glitch in the electrical recharging phase of the heart.

What is the QT interval?

The QT interval is a time interval on the ECG. It represents the time from the electrical stimulation (depolarization) of the heart's pumping chambers (ventricles), to the end of the recharging of the electrical system (repolarization). It is measured in milliseconds and closely approximates the time from the beginning of the ventricles' contraction until the end of relaxation.

Patients who continue to have symptoms in spite of appropriate doses of beta-blockers, or are not able to tolerate their beta-blocker therapy, may also require additional medications, surgery (left cardiac sympathetic denervation [LCSD]), or device therapy with an implantable cardioverter defibrillator (ICD). Patients who have experienced a cardiac arrest usually receive an ICD. With LQTS genetic testing, treatment strategies can be guided by the underlying genetic cause. For example, beta-blockers are generally extremely protective in type 1 LQTS (LQT1) and are also protective for the majority of patients with type 2 and type 3 LQTS (LQT2 and LQT3). LCSD surgery is available in a few LQTS centers providing another important treatment option for those patients who continue to have syncope despite beta-blockers, or have experienced appropriate ICD shocks or are not tolerating their medications because of excessive side effects.

Persons over 40 years of age at the time of diagnosis (perhaps > 20 for LQT1 males) who have been asymptomatic (without symptoms) all their life (or for many, many years) may not need any active treatment, as their risk of developing symptoms at later ages is very low. As with all patients with LQTS, these seemingly low-risk older patients need to avoid low blood potassium (caused by diuretic drug use, vomiting or diarrhea) and drugs which aggravate the heart's recharging system and prolong the QT interval. This combination of an otherwise dormant LQTS genetic substrate plus a 'second hit' can provide a fatal 1-2 punch. For a complete list of drugs that prolong the QT interval and/or induce torsade de pointes, visit www.StopSADS.org or www.crediblemeds.org.

Medication compliance

It is very important that LQTS-directed medical therapies be taken every day and not missed or omitted. The medications are not curative; they only provide protection while being taken and depending on the beta blocker, the protective effect can disappear within a day. After that the risk of cardiac events is the same as if the patient had not taken the medication at all. In addition, dose skipping/missing can actually be dangerous as the heart can become hyperexcitable when the medicine is not present. Parents should teach their children about the importance of daily medication, and should make sure each daily dose is taken. Physicians need to discuss this directly with all patients, but particularly pre-teens and teenagers. **The most common reason for cardiac events while on medication may be that the medication has been missed or stopped.**

How can parents protect their kids?

- Make sure children take their medication daily, no missing doses.
- See the doctor regularly for follow-up. Growing children need medication dose changes regularly. Make sure you see the doctor at least once a year, more frequently during very rapid growth, and discuss the need for dose changes. These drugs are given according to body weight; so, in growing children dose changes may be frequent.
- If your child has a passion for sports, your child should be evaluated by a physician with expertise in LQTS to determine if continuation in competitive sports is reasonable.
- Get additional medical advice if you are not comfortable with how things are going. Ideally, every patient/family with LQTS should be cared for by a heart rhythm specialist (cardiac electrophysiologist) or even a long QT syndrome specialist. Do not hesitate to obtain a second opinion if you have any questions about your or your child's treatment.
- Learn CPR, encourage your child's school district to implement an automatic external defibrillator (AED) in their schools, and consider obtaining a personal AED as part of the family's LQTS safety gear.
- It is important to be sure that your child's school is aware of his or her condition and is properly prepared to respond in the event of an emergency. Visit www.StopSADS.org for care plans and other important materials that will help you in meeting with school personnel to ensure a safe environment for your child.



SADS Foundation
508 East South Temple, Suite 202
Salt Lake City, UT 84102
800-STOP SAD
www.StopSADS.org
sads@sads.org

Updated 8/2014 Reviewers: Michael Ackerman, MD, Peter Schwartz, MD, Arthur Wilde, MD

The Sudden Arrhythmia Death Syndromes (SADS) Foundation is a leader in education, research and advocacy. Our Mission is to save the lives and support the families of children & young adults who are genetically predisposed to sudden death due to heart rhythm abnormalities.


www.StopSADS.org



The QT interval varies in each person and between persons like most physiologic parameters, such as blood pressure or heart rate. In particular, the QT varies with the heart rate. It shortens as the rate increases and lengthens as the rate decreases. Therefore, there is a range of normal or healthy QT intervals. In contrast, the long QT heart often recharges sluggishly or inefficiently as evidenced by a “prolonged QT interval” on the ECG.

To determine if a given QT is normal for a given heart rate, the QT is corrected for the heart rate using a simple mathematical formula, and the resultant quantity is called the QTc. The QTc is the value that doctors generally use when assessing for LQTS.

The average QTc value is around 400 ± 20 ms. Probably less than 1% of the population has a QTc < 350 ms and less than 1% has a QTc > 480 ms. About 90% of people have a value between 380 ms and 440 ms, which doctors generally consider as the “normal” range. However, this “normal” range is affected by age, particularly relevant in children, and gender. For example, women tend to have slightly longer QT intervals than men. While a QTc of 440 ms represents the top 2.5 percentile in a 4-day old infant, this same QTc value is seen in 10 – 20% of postpubertal women. The diagram below provides an example of a normal QTc and a prolonged QTc.



What are the symptoms of LQTS?

Nearly half of patients with LQTS **NEVER** have a symptom. However, if/when the LQTS heart “spins electrically out of control” in its trademark cardiac arrhythmia (called torsade de pointes) sudden, temporary, loss of consciousness (syncope) is the most common event. These events usually occur without warning and are often triggered by exertion, emotions, or auditory stimuli. Typically, the onset of symptoms is earlier in boys than in girls. Statistically, the greatest risk

window includes the first three decades of life (the first two decades in males and the second and third decade in females). However, there are tragic exceptions to these trends and LQTS-related events can continue in people in their 40s and 50s. When presenting later, a second hit, due to a medication that further aggravates the QT interval or low potassium, is often present.

In patients who experience syncope only, the torsade de pointes rhythm spontaneously returns to normal, usually in less than a minute, and the patient quickly regains consciousness, without disorientation or confusion. Some patients may experience slight fatigue afterwards; others feel fine and resume their regular activities. If this bad LQTS rhythm persists longer, patients may then manifest a generalized seizure. In fact, some patients with LQTS have been misdiagnosed initially with epilepsy and even treated with anti-epileptic medication. In both the syncope and seizure presentations, the long QT heart reverted back to normal sinus rhythm, and the “spell” is over. On the other hand, in a minority of patients, the torsade de pointes rhythm persists longer still and degenerates further into the heart rhythm known as ventricular fibrillation, which rarely reverts back to a normal rhythm without medical intervention. If the ventricular fibrillation is not (immediately) converted, usually by electrical defibrillation, the outcome is sudden cardiac death or sudden cardiac arrest.

When should the diagnosis be suspected?

- In any young person with **unexplained syncope (fainting), unexplained seizures, or unexplained cardiac arrest or sudden death**. Usually, a careful history of the events surrounding the syncope differentiates LQTS-induced syncope from the common faint, known as vasovagal or neurocardiogenic syncope. The LQTS faint is usually precipitous and without warning. It often occurs during or just after **physical exertion (including swimming), emotional excitement including fear or sudden auditory arousal** (such as a doorbell or alarm clock), but may occur during sleep or at rest. Conversely, in vasovagal syncope, most times there are warning symptoms such as dizziness, blurring or blackening of vision, tingling or sweating, for seconds to even minutes prior to the syncope. Also, a precipitating event is usually present, commonly pain, injury, nausea, or an unpleasant or stressful experience.



- When there is a **family history** of unexplained syncope, unexplained seizures, or sudden death in young people. As noted above, at least one-third to one-half of individuals never exhibit symptoms, and, therefore, the lack of prior symptoms does not exclude a person or family from having LQTS.
- When the autopsy is normal following the sudden and unexpected death of a young person.

How is the diagnosis made?

LQTS is suspected based upon the patient’s personal story, his/her family story, and careful examination of the ECG. If the story is suspicious, a QTc exceeding 470 ms in males and 480 ms in females is sufficient evidence for a diagnosis of probable LQTS, assuming that medications which prolong the QT interval or other QT-prolonging medical conditions have been ruled out. Importantly, as an incidental ECG finding, QTc values exceeding these cut-off values do NOT equal the diagnosis of LQTS. On the other hand, a QTc of less than 400 ms in males and 410 ms in females makes the diagnosis unlikely. **The computer generated QTc may be incorrect, so when the diagnosis of LQTS is considered, the physician must confirm manually the QTc measurement.**

However, not all LQTS patients have a prolonged QTc on the initial ECG. In fact, about 30 – 50% have a QTc that overlaps with the normal range and at least 25% of affected relatives have a QTc < 440 ms. QTcs in this range are inconclusive, and therefore must be clarified through additional testing that may include exercise stress testing, the epinephrine QT stress test, and genetic testing.

How is LQTS inherited, and who in a known or suspected family should be tested?

LQTS is usually inherited by autosomal dominant transmission. This means that it generally affects boys and girls equally, and that each child of an affected parent has a 50% chance of inheriting the genetic abnormality. In a really large family, close to 50% of the children would inherit the LQTS-causing genetic abnormality. In average size families, it can range from all to none as each child has an independent 50/50 chance of inheriting the particular disease gene. Once a family member is identified with LQTS, it is extremely important that other family members be tested for the syndrome. It is especially important to know which parent and grandparent has the abnormality, since brothers and sisters, aunts, uncles, nephews, nieces, and cousins on the affected side are potentially at risk.

This prospective screening, by ECG, is extremely important so that all affected family members are identified and treated early in order to prevent the tragic and unnecessary sudden deaths that may otherwise occur.

What about genetic testing?

Genetic testing for LQTS is now the “standard of care” and it is increasingly being used to diagnose and treat people. For people with very prolonged QT intervals, the genetic test will help your physician recommend treatment. It will also help your family members to get diagnosed and treated. For people with a “borderline” diagnosis because the QTc is almost normal and they have no symptoms, genetic testing can help “rule out” or “confirm” the diagnosis of LQTS.

Your physician will order the initial test on one family member (so-called index case). Once/if a family member has a gene mutation identified, testing of other family members for that specific mutation is available, and can assist in clarifying those family members with non-definitive ECG findings. In fact, if a genetic diagnosis of LQTS is established for the index case, the **ONLY** definitive test to rule in or rule out LQTS for family members and relatives is the LQTS genetic test. It is also very reassuring to know that certain members of the family are not affected and that, therefore, they will not transmit the disease to their offspring.

There are a number of commercial, CLIA-certified labs from which you can order your genetic testing. As with any health-care service, it is always a good idea to be a wise consumer and research each company to see which one might be the better choice for you. Visit www.StopSADS.org for more information.

What is the treatment and who should be treated?

All **symptomatic** patients should receive treatment. Most children and young adults should be treated even if they do not have symptoms. This is because symptoms might occur and sudden death may be the first symptom. At present, it is not possible to tell which child or youth is destined to have symptoms. Thus, preventative treatment is recommended for most. The usual treatment involves taking beta-blocker medications daily. This approach is effective for the majority of patients with LQTS. The dose needs to be monitored closely, balancing the prevention of LQTS spells with unwanted side effects related to energy level and mood.